



***Finerenone in Heart Failure and Chronic Kidney Disease with Type 2 Diabetes: the FINE-HEART Pooled Analysis of Cardiovascular, Kidney, and Mortality Outcomes***

**Muthiah Vaduganathan on behalf of**


**Gerasimos Filippatos; Brian Claggett; Akshay Desai; Pardeep Jhund; Alasdair Henderson; Meike Brinker; Peter Kolkhof; Patrick Schloemer; James Lay-Flurrie; Prabhakar Viswanathan; Carolyn Lam; Michele Senni; Sanjiv Shah; Adriaan A. Voors; Faiez Zannad; Peter Rossing; Luis Ruilope; Stefan Anker; Bertram Pitt; Rajiv Agarwal; John McMurray; Scott Solomon**

**PROSPERO CRD42024570467**




# Strong Epidemiological Overlap of Cardiovascular, Metabolic, and Kidney Disorders

US NHANES survey cycles 1999–2020



US NHANES Survey Cycles 1999-2020  
Ostrominski J...Vaduganathan M. JAMA Cardiology 2023

US NHANES survey cycles 2011–2020




US NHANES Survey Cycles 2011-2020  
Aggarwal R...Vaduganathan M. JAMA 2024



# Could the Non-Steroidal MRA, Finerenone, Modify Risk across the Cardio-Kidney-Metabolic Spectrum?

- Finerenone is a non-steroidal MRA that has been studied in RCTs of patients with T2D and CKD and separately in patients with HF (with and without T2D).
- However, none of these trials were individually powered to evaluate treatment effects on mortality outcomes or effects in key subgroups.



# Design of FINE-HEART Umbrella Program



Prospectively Registered:  
PROSPERO CRD42024570467

(n=18,991 Participants)






Pooling data in the FINE-HEART program increased precision to robustly assess the efficacy and safety of the non-steroidal MRA finerenone on important cardio-kidney outcomes and is enriched for participants with a high burden of CKM multimorbidity.

## Study Designs of the Individual Trials

	<b>FINEARTS-HF</b>	<b>FIDELIO-DKD and FIGARO-DKD</b>
<b>Validly Randomized</b>	6,001	12,990
<b>Countries</b>	37	48
<b>Patient population</b>	HFmrEF or HFpEF	CKD and T2D
<b>Inclusion criteria</b>	<ul style="list-style-type: none"> <li>• Adults (<math>\geq 40</math> years)</li> <li>• Symptomatic HF</li> <li>• LVEF <math>\geq 40\%</math></li> <li>• Elevation natriuretic peptides</li> <li>• Structural heart disease</li> <li>• Recent diuretic use</li> </ul>	<ul style="list-style-type: none"> <li>• Adults (<math>\geq 18</math> years old)</li> <li>• T2D</li> <li>• UACR <math>\geq 30</math> mg/g</li> <li>• Maximally tolerated RASi</li> </ul>
<b>Exclusion criteria</b>	Potassium $\leq 5.0$ mmol/L	Potassium $\leq 4.8$ mmol/L
<b>Dosage and titration</b>	eGFR $\leq 60$ : 10 up to 20 mg eGFR $> 60$ : 20 up to 40 mg (potentially down to 10 mg)	eGFR $< 60$ : 10 up to 20 mg eGFR $\geq 60$ : 20 mg (potentially down to 10 mg)
<b>Study duration</b>	2.6 years	2.6 years (FIDELIO-DKD) 3.4 years (FIGARO-DKD)

# Baseline Characteristics of FINE-HEART Integrated Population


	Finerenone (n=9,501)	Placebo (n=9,490)
<b>Age</b>	<b>67±10</b>	<b>67±10</b>
<b>Women</b>	<b>36%</b>	<b>35%</b>
<b>White Race</b>	<b>72%</b>	<b>72%</b>
<b>BMI (kg/m<sup>2</sup>)</b>	<b>31±6</b>	<b>31±6</b>
<b>Systolic BP (mmHg)</b>	<b>135±15</b>	<b>134±15</b>
<b>Potassium (mmol/L)</b>	<b>4.4±0.5</b>	<b>4.4±0.5</b>
<b>eGFR (mL/min/1.73m<sup>2</sup>)</b>	<b>59±21</b>	<b>59±21</b>
<b>&lt;25</b>	<b>1%</b>	<b>1%</b>
<b>25 to &lt;45</b>	<b>29%</b>	<b>29%</b>
<b>45 to &lt;60</b>	<b>27%</b>	<b>26%</b>
<b>≥60</b>	<b>44%</b>	<b>44%</b>
<b>UACR (mg/g)</b>	<b>283</b> <b>[46-836]</b>	<b>293</b> <b>[47-855]</b>
<b>A1: &lt;30</b>	<b>20%</b>	<b>20%</b>
<b>A2: 30 to &lt;300</b>	<b>31%</b>	<b>31%</b>
<b>A3: ≥300</b>	<b>49%</b>	<b>50%</b>




	Finerenone (n=9,501)	Placebo (n=9,490)
<b>HbA1c (%)</b>	<b>7.3±1.4</b>	<b>7.3±1.4</b>
<b>HF</b>	<b>37%</b>	<b>37%</b>
<b>Diabetes</b>	<b>81%</b>	<b>81%</b>
<b>CKD</b>	<b>84%</b>	<b>84%</b>
<b>AF</b>	<b>15%</b>	<b>15%</b>
<b>Diuretics</b>	<b>66%</b>	<b>67%</b>
<b>ACEi/ARB/ARNI</b>	<b>93%</b>	<b>93%</b>
<b>Statins</b>	<b>70%</b>	<b>71%</b>
<b>SGLT2i</b>	<b>9%</b>	<b>9%</b>
<b>GLP-1RA</b>	<b>6%</b>	<b>6%</b>

# High Burden of Cardio-Kidney-Metabolic Disease Overlap


## Baseline CKM Status in FINEHEART



# Primary Endpoint: CV Death




**Primary Analysis:**  
**CV Death Excluding Unknown Deaths**  
**Finerenone 421 (4.4%) vs. Placebo 471 (5.0%)**




**Prespecified Sensitivity Analysis:**  
**CV Deaths Including Unknown Deaths**  
**Finerenone 627 (6.6%) vs. Placebo 703 (7.4%)**



# Secondary Endpoint: All-Cause Mortality




# Secondary Endpoint: HF Hospitalization



# Secondary Endpoint: Kidney Composite Endpoint

sustained eGFR decline of  $\geq 50\%$ , kidney failure\*, or death due to kidney failure


## Kidney Composite Endpoint





\*sustained eGFR < 15 ml/min/1.73m<sup>2</sup>, chronic dialysis, or kidney transplantation

# Summary of Prespecified Efficacy Endpoints

Outcome		HR (95% CI)	P-value
<b>Primary Endpoint</b>			
CV death (excluding undetermined death)		0.89 (0.78–1.01)	0.076
<i>Prespecified sensitivity analysis:</i> CV death (including undetermined death)		0.88 (0.79–0.98)	0.025
<b>Secondary Endpoints</b>			
Kidney Composite Endpoint		0.80 (0.72–0.90)	<0.001
HF Hospitalization		0.83 (0.75–0.92)	<0.001
CV Death or HF Hospitalization		0.85 (0.78–0.93)	<0.001
New-onset Atrial Fibrillation		0.83 (0.71–0.97)	0.018
Major Adverse Cardiovascular Events		0.91 (0.85–0.98)	0.010
All-cause Death		0.91 (0.84–0.99)	0.027
All-cause Hospitalization		0.95 (0.91–0.99)	0.025
All-cause Death or All-cause Hospitalization		0.94 (0.91–0.98)	0.007



# Broad Consistency Across 17 Prespecified Subgroups for the Primary Endpoint (CV Death)



## Safety Outcomes

	<b>Finerenone</b>	<b>Placebo</b>
	n=9,482	n=9,467
<b>Any serious adverse event</b>	35%	37%
<b>Any ae leading to treatment discontinuation</b>	5%	5%
<b>Any potassium &gt;5.5 mmol/L</b>	17%	8%
<b>Any potassium &gt;6.0 mmol/L</b>	3%	1%
<b>Any potassium &lt;3.5 mmol/L</b>	5%	10%
<b>Hyperkalemia</b>	13%	6%
<b>Hyperkalemia leading to discontinuation</b>	1.3%	0.5%
<b>Hyperkalemia leading to hospitalization</b>	0.8%	0.2%
<b>Hyperkalemia leading to death</b>	0%	0%
<b>Acute kidney injury</b>	4%	3%
<b>Acute kidney injury leading to discontinuation</b>	0.2%	0.1%
<b>Acute kidney injury leading to hospitalization</b>	2%	1%
<b>Systolic blood pressure&lt;100mmHg</b>	11%	7%
<b>Gynecomastia or breast hyperplasia</b>	0.2%	0.2%

Treatment-emergent adverse events are defined as any adverse event occurring in any patient who has received at least one dose of study drug and within 3 days of permanent discontinuation. This safety table includes 1 patient who was randomized to placebo but who actually received finerenone.


## Conclusions

- The FINE-HEART participant-level pooled analysis represents the largest analysis of the effects of the non-steroidal MRA finerenone across the CKM spectrum.
- While this pooled analysis failed to demonstrate significant reductions in cardiovascular death, finerenone was associated with significantly lower deaths of any cause, cardiovascular events, and kidney outcomes.
- Treatment effects were consistent across all tested clinical subgroups including those with multiple intersecting CKM conditions and on background SGLT2i or GLP-1RA.
- No new or unexpected safety signals were uncovered in this pooled analysis.

**The totality of the evidence supports the disease-modifying potential of finerenone in broad, high-risk patient populations encompassing cardiovascular, kidney, and metabolic diseases.**

Full Details Available Online in *Nature Medicine*

nature  
medicine



**Finerenone in Heart Failure and Chronic Kidney Disease with Type 2 Diabetes: the FINE-HEART Pooled Analysis of Cardiovascular, Kidney, and Mortality Outcomes**

Muthiah Vaduganathan; Gerasimos Filippatos; Brian L. Claggett; Akshay S. Desai; Pardeep S. Jhund; Alasdair Henderson; Meike Brinker; Peter Kolkhof; Patrick Schloemer; James Lay-Flurrie; Prabhakar Viswanathan; Carolyn SP Lam; Michele Senni; Sanjiv J Shah; Adriaan A. Voors; Faiez Zannad; Peter Rossing; Luis M. Ruilope; Stefan D. Anker; Bertram Pitt; Rajiv Agarwal; John JV McMurray; Scott D. Solomon

<https://doi.org/10.1038/s41591-024-03264-4>



# In Memory of the Late Dr. George Bakris (1952-2024)



A pioneer in cardio-kidney-metabolic research,  
physician, leader, colleague, and dear friend